

CPDO Examination - Sample Solo Item 1

A 7-year-old patient has acute streptococcal erysipelas involving the eyelids. Which of the following is the best oral medication for this patient?

(Item type - Treatment/Management)

- a. Penicillin V
- b. Tetracycline
- c. Doxycycline
- d. Cyclosporine

405839 Rev 10.28.13

CPDO Examination - Sample Solo Item 2

A patient has a mucopurulent discharge from a lesion of the right lower eyelid. Which of the following ophthalmic ointments would be most effective to use if a Gram stain of the discharge reveals predominantly polymorphonuclear leukocytes and Gram-negative organisms?

(Item type - Clinical Correlation of Basic Science Principles)

- a. Ciprofloxacin
- b. Fluorometholone
- c. Erythromycin
- d. Sulfacetamide

405750 Rev 10.28.13

CPDO Examination - Sample Solo Item 3

A 78-year-old white female tells you that the vision in her right eye gradually has been worsening for the past 2 months. You find that her best corrected visual acuities are 20/60 OD and 20/20 OS. Ophthalmoscopy and macular OCT reveal hemorrhage due to a subfoveal choroidal neovascular membrane OD. Of the following, which is the best management for this patient's right eye?

(Item type - Treatment/Management)

- a. Intravitreal bevacizumab
- b. Direct argon laser photocoagulation
- c. Prednisone
- d. Follow-up retinal examination in 3 months

405657 Rev 10.28.13

CPDO Examination - Sample Solo Item 4

**You place a 21-year-old female on oral tetracycline 250 mg po qid for 30 days. This long-term treatment may cause all of the following side effects except:
(Item type - Related to Treatment/Management)**

- a. skin photosensitivity.
- b. papilledema.
- c. gastrointestinal distress.
- d. ototoxicity.

405818 Rev 10.28.13

CPDO Examination - Sample Solo Item 5

**Which 2 of the following are most appropriate in the initial management of an adult male who is experiencing an acute anaphylactic reaction? (Select 2 answer options)
(Item type - Treatment/Management)**

- a. Expose the patient to ammonia inhalants.
- b. Administer systemic epinephrine.
- c. Give the patient orange juice.
- d. Have the patient breathe into a paper bag.
- e. Instruct a bystander to call 911 immediately.
- f. Perform the Heimlich maneuver.

405928 Rev 10.28.13